



## Tsunami early warning using earthquake rupture duration

Anthony Lomax<sup>1</sup> and Alberto Michelini<sup>2</sup>

Received 9 January 2009; revised 28 February 2009; accepted 27 March 2009; published 9 May 2009.

[1] Effective tsunami early warning for coastlines near a tsunamigenic earthquake requires notification within 5–15 minutes. We have shown recently that tsunamigenic earthquakes have an apparent rupture duration,  $T_0$ , greater than about 50 s. Here we show that  $T_0$  gives more information on tsunami importance than moment magnitude,  $M_w$ , and we introduce a procedure using seismograms recorded near an earthquake to rapidly determine if  $T_0$  is likely to exceed  $T = 50$  or 100 s. We show that this “duration-exceedance” procedure can be completed within 3–10 min after the earthquake occurs, depending on station density, and that it correctly identifies most recent earthquakes which produced large or devastating tsunamis. This identification forms a complement to initial estimates of the location, depth and magnitude of an earthquake to improve the reliability of tsunami early warning, and, in some cases, may make possible such warning. **Citation:** Lomax, A., and A. Michelini (2009), Tsunami early warning using earthquake rupture duration, *Geophys. Res. Lett.*, 36, L09306, doi:10.1029/2009GL037223.

### 1. Introduction

[2] Effective tsunami early warning for coastlines near a tsunamigenic earthquake requires notification within 5–15 minutes after the earthquake origin time (OT). Organizations such as the Japan Meteorological Agency (JMA), the German-Indonesian tsunami early warning system (GITEWS) and the West Coast and Alaska (WCATWC), and Pacific (PTWC) Tsunami Warning Centers first identify potentially tsunamigenic earthquakes based on rapidly determined earthquake parameters such as location, depth and magnitude. JMA issues warnings for Japan about 3 min after OT for events expected to produce a tsunami with height exceeding 0.5 m. GITEWS issues warnings for Indonesia within 5 min after OT based on the earthquake parameters and corresponding, pre-calculated tsunami scenarios. WCATWC and PTWC issue regional warning notifications within about 5–10 min after OT for shallow, underwater events around North America and in the Pacific basin with moment magnitude  $M_w \geq 7.5$  [e.g., *Hirshorn and Weinstein, 2009*].

[3] Recently, through analysis of teleseismic,  $P$ -wave seismograms ( $30^\circ$ – $90^\circ$  great-circle distance; GCD), we have shown that an apparent rupture duration,  $T_0$ , greater than about 50 s forms a reliable indicator for tsunamigenic earthquakes [Lomax and Michelini, 2009] (LM2009). Here we exploit this result and introduce a “duration-exceedance”

procedure to rapidly determine if  $T_0$  for an earthquake is likely to exceed 50 or 100 s and thus to be a potentially tsunamigenic earthquake. This procedure does not require accurate knowledge of the earthquake location or magnitude and can be completed within 5–10 min after OT for most regions in the world.

### 2. Tsunami Importance, Moment Magnitude, and Rupture Duration


[4] We consider a reference set of 76 underwater earthquakes since 1992 with  $M_w \geq 6.6$  (Table S1<sup>1</sup>). Since there is currently no uniform, physical measure of size available for most tsunamis, following LM2009, we define an approximate measure of tsunami importance,  $I_t$ , based on 0–4 descriptive indices,  $i$ , of tsunami effects (deaths, injuries, damage, houses destroyed), and maximum water height  $h$  in meters from the NOAA/WDC Historical Tsunami Database ([http://www.ngdc.noaa.gov/hazard/tsu\\_db.shtml](http://www.ngdc.noaa.gov/hazard/tsu_db.shtml)):  $I_t = i_{height} + i_{deaths} + i_{injuries} + i_{damage} + i_{houses-destroyed}$ , where  $i_{height} = 4, 3, 2, 1, 0$  for  $h \geq 10, 3, 0.5$  m,  $h > 0$  m,  $h = 0$  m respectively. We set  $I_t = 0$  for events not in the database, and note that  $I_t$  is approximate and unstable since it depends strongly on the available instrumentation, coastal bathymetry and population density in the event region.  $I_t \geq 2$  corresponds approximately to the JMA threshold for issuing a “Tsunami Warning”; the largest or most devastating tsunamis typically have  $I_t \geq 10$ .

[5] Figure 1 shows a comparison of  $I_t$  with the Global Centroid-Moment Tensor (CMT) moment-magnitude,  $M_w^{CMT}$  [Dziewonski et al., 1981; Ekström et al., 2005], and with  $T_0$  durations calculated from high-frequency,  $P$ -wave seismograms at teleseismic distance following the procedure of LM2009. The thresholds  $M_w^{CMT} \geq 7.5$  and  $T_0 \geq 50$  s both identify most of the events with  $I_t \geq 2$  (see also Tables 1 and S1).  $M_w^{CMT}$ , however, shows no clear relationship to  $I_t$  or to event type; in contrast,  $T_0$  tends to increase for larger  $I_t$ , especially for tsunami earthquakes (type T; characterized by unusually large tsunamis and a deficiency in moment release at high frequencies) [e.g., *Satake, 2002*]. We do not consider here the energy-to-moment parameter,  $\Theta$ , which is useful for identification of tsunami earthquakes [Newman and Okal, 1998], because it is not a good indicator for tsunamigenic events in general (e.g., LM2009).

[6] Since CMT-based  $M_w$  magnitudes are only available 30 min or later after OT, rapid magnitude estimates such as  $M_{wp}$  [Tsuboi et al., 1995, 1999] are used for tsunami warning. But  $M_{wp}$  performs poorly relative to  $M_w^{CMT}$  or  $T_0$  for identifying events with  $I_t \geq 2$  (Table 1). Other rapid magnitude estimates for large earthquakes (e.g., *Hara [2007]*, LM2009,  $M_{wpd}$ ; and *Bormann and Saul [2009]*,  $m_{Bc}$ ) may perform nearly as well as  $M_w^{CMT}$  or  $T_0$  (e.g.,  $M_{wpd}$

<sup>1</sup>ALomax Scientific, Mouans-Sartoux, France.

<sup>2</sup>Istituto Nazionale di Geofisica e Vulcanologia, Rome, Italy.



**Figure 1.** Comparison of tsunami importance  $I_t$  with (top) moment-magnitude  $M_w^{CMT}$  and (bottom) apparent source duration,  $T_0$ , calculated from teleseismic observations. Event labels show event type for non interplate-thrust events with  $I_t \geq 2$ .  $T$ , tsunami earthquake;  $P$ , intraplate;  $So$ , strike-slip oceanic;  $S$ , strike-slip continental; and  $R$ , reverse-faulting.

in Tables 1 and S1), but are not available until about 15 min or later after OT. Thus very rapid determination of a large  $T_0$ , e.g.  $T_0 \geq 50$  s, would provide important complementary information to initial location, depth and magnitude estimates for early assessment of earthquake tsunamigenic potential.

### 3. Methodology for Rapid Rupture Duration Determination

[7] We determine if  $T_0$  for an earthquake is likely to exceed pre-determined thresholds  $T = 50, 100$  s through high-frequency (HF) analysis of vertical-component, broadband seismograms [e.g., Lomax, 2005; Lomax and Michelini, 2005; Lomax et al., 2007; LM2009]. We proceed

as follows for each seismogram (Figure 2): 1) apply a 4-pole, 1–5 Hz Butterworth band-pass filter to form a HF trace; 2) auto-pick the  $P$  arrival time on the HF trace; 3) measure  $A_{ref}$ , the *rms* amplitude for the first 25 s after the  $P$  time on the HF trace; 4) calculate the ratio of the *rms* HF amplitude from 50–60 s after the  $P$  time with  $A_{ref}$  to obtain a station duration-exceedance level for 50 s,  $l_{50}$ , and a similar ratio for 100–120 s after  $P$  with  $A_{ref}$  to obtain  $l_{100}$ .

[8] We define event duration-exceedance levels,  $L_T$ ,  $T = 50, 100$  s, as the median (50 percentile) of the station  $l_{50}$ ,  $l_{100}$  values after removing the upper 10 percentile of values to avoid noisy or anomalously long HF signals. If an event exceedance level  $L_T$  is greater (less) than 1.0, then  $T_0$  is likely (unlikely) to exceed  $T$  seconds. This procedure does not require an event location or magnitude, and all processing can be performed in the time domain; indeed, individual station  $l_{50}$  and  $l_{100}$  values can be calculated autonomously at each station.

### 4. Application to Reference Earthquakes

[9] We apply the duration-exceedance procedure to the reference earthquakes using data up to 10 min after OT from stations at 0–30° GCD from each event to simulate the information available in the first minutes after an earthquake occurs. The  $L_{50}$  exceedance level results are tabulated in Table 1 and all event parameters and exceedance level results are tabulated in Table S1; plots of the time evolution of the  $L_{50}$  calculation for two events are shown in Figure 3 and for  $L_{50}$  and  $L_{100}$  for selected events are shown in Figure S1.

[10] A comparison of  $L_T$ ,  $T = 50, 100$  s, with the  $T_0$  durations calculated from teleseismic observations (Figure 4 (top) and Table S1) shows that, in general, the duration-exceedance level  $L_T$  increases with increasing  $T_0$  and is greater than 1 for events with  $T_0 > T$ . There is much scatter in these results, due primarily to the difficulty in determining cutoff points on the HF seismograms (e.g., Figure 2 and LM2009), but they confirm that the rapidly available  $L_T$  measures form reliable proxies for the teleseismic,  $T_0$  durations.

### 5. Discussion


[11] A comparison of the  $L_{50}$  exceedance level with tsunami importance,  $I_t$ , (Figure 4 (bottom) and Tables 1 and S1) shows correct identification ( $L_{50} \geq 1$ ) of most events with  $I_t \geq 2$ . The miss-identified events are a shallow, offshore thrust event,  $I_t = 8$ , 2003.05.21,  $M_w$ 6.8, N Algeria, and two shallow, oceanic, strike-slip events,  $I_t = 13$ , 1994.11.14,  $M_w$ 7.1, Philippines and  $I_t = 9$ , 2006.03.14,  $M_w$ 6.7, Seram Indonesia. All of these events are also missed

**Table 1.** Results for  $L_{50}$  Classification of tsunamigenic Earthquakes<sup>a</sup>

Discriminant	Available (min after OT)	Critical Value	Correctly Identified			Missed		False	
			$I_t \geq 2$	Percentage <sup>b</sup>	$I_t < 2$	$I_t \geq 2$	$I_t < 2$		
$M_w^{CMT}$	30+	7.5	27	87	34	4	11		
$T_0$ (teleseismic)	15+	50	26	84	32	5	13		
$M_{wpd}$ (raw)	15+	7.5	24	77	32	7	12		
$M_{wp}$	3–10	7.5	16	52	38	15	7		
$L_{50}$	3–10	1.0	28	90	32	3	13		

<sup>a</sup>76 events classified; 31 have  $I_t \geq 2$ .

<sup>b</sup>Percentage of all events with  $I_t \geq 2$  that are correctly identified.



**Figure 2.** Raw, broadband velocity seismogram, HF seismogram and smoothed  $rms$  amplitude of HF seismogram for two events: (top) 2006.07.17,  $M_w 7.7$ ,  $T_0 = 180$  s, and  $I_t = 18$  Indonesia tsunami earthquake recorded at station COCO at  $11^\circ$  GCD and (bottom) 2008.04.09,  $M_w 7.0$ ,  $T_0 = 23$  s, and  $I_t = 0$  Loyalty Islands interplate thrust recorded at station AFI at  $19^\circ$  GCD. OT, origin time; P, automatic  $P$  pick; P to Ar, T50 and T100, time windows (shaded) for calculation of  $rms$  HF amplitude for  $A_{ref}$ ,  $l_{50}$  and  $l_{100}$ , respectively.

using the magnitude discriminant,  $M_w \geq 7.5$ , and thus produced larger than expected tsunamis. There are 13 events with  $I_t < 2$  that are falsely identified by  $L_{50} \geq 1$  values as likely tsunamigenic ( $I_t \geq 2$ ); 7 of these events have  $I_t = 1$  and thus produced small tsunamis, while some may have involved under land or strike-slip rupture, or produced unobserved tsunamis. The remaining events with  $I_t < 2$  are correctly identified as unlikely tsunamigenic by  $L_{50} < 1$  values. For most events, the  $L_{50}$  values have stabilized within 4–6 min after OT (Figures 3 and S1).

[12] The  $L_{50}$  discriminant correctly identifies 90% of tsunamigenic events with  $I_t \geq 2$ . The overall performance of the  $L_{50}$  discriminant is similar to that of  $M_w^{CMT}$ ,  $M_{wpd}$ , and teleseismic  $T_0$  (Table 1), though these latter three measures are not available until at least 30, 15 and 15 min, respectively, after OT (LM2009). In contrast, the rapidly available  $M_{wp}$  discriminant correctly identifies only 52% of tsunamigenic events with  $I_t \geq 2$ , primarily because  $M_{wp}$  under-


estimates the size of events with  $M_w^{CMT} > 7.0$ –7.5, particularly tsunami earthquakes and other events with long rupture duration (e.g., LM2009).

[13] The results for  $L_{100}$  (Figure 4 and Table S1) show that  $L_{100} \geq 1$  identifies well events with longer duration,  $T_0$ , events with  $I_t \geq 10$ , and most tsunami earthquakes (type T). In contrast, 1994.11.14 Philippines, 1998.07.17 Papua New Guinea, and two intraplate events (type P) with only moderately long  $T_0$  but large  $I_t$  have  $L_{100} < 1$  values. For events in regions with denser station coverage, the  $L_{100}$  values have stabilized by 6–8 min after OT (Figure S1).


[14] Since the station  $l_T$  exceedance values can be calculated autonomously at each station, they could aid in providing very early, local tsunami warning. For example, the first station  $l_{50}$  values for the 2006 Indonesian event in Figure 3 are available only 2–4 min after OT. Single  $l_T$  exceedance values must be used with care, however, as they can be biased at small epicentral distances by HF radiation effects and secondary phases, especially  $S$ .

## 6. Conclusions

[15] We have shown that apparent rupture duration,  $T_0$ , provides more information on tsunami importance,  $I_t$ , than



**Figure 3.** Evolution for 10 min after OT of the  $T_0 > 50$  s exceedance level ( $L_{50}$ ) calculation for (top) 2006.07.17,  $M_w 7.7$ ,  $T_0 = 180$  s, and  $I_t = 18$  Indonesia tsunami earthquake and (bottom) 2008.04.09,  $M_w 7.0$ ,  $T_0 = 23$  s, and  $I_t = 0$  Loyalty Islands interplate thrust. Blue lines show  $P$ -arrival times for each station; red, yellow or green horizontal bars show the station exceedance levels,  $l_{50}$ , starting at its first reported time (about 60 s after the corresponding  $P$  time). Histogram shows  $l_{50}$  values at 600s; the median (50 percentile) and bounds (20 and 80 percentile), respectively, for  $L_{50}$  are indicated by solid and dotted white lines on the main plot and as a colored diamond and error bar. Red indicates  $l_{50}$ (or  $L_{50}$ )  $\geq 1$  (likely that  $T_0 > 50$  s and  $I_t \geq 2$ ); yellow indicates  $0.7 \leq l_{50}$ (or  $L_{50}) < 1$  (possible that  $T_0 > 50$  s and  $I_t \geq 2$ ); and green indicates  $l_{50}$ (or  $L_{50}) \leq 0.7$  (unlikely that  $T_0 > 50$  s or  $I_t \geq 2$ ). For both events the  $L_{50}$  values have stabilized by 4–6 min after OT. For real-time monitoring, comprehensive information about exceedance level could be provided by a time-sliding display similar to the above.



**Figure 4.** Comparison of exceedance levels  $L_{50}$  and  $L_{100}$  with (top) apparent source duration  $T_0$  calculated from teleseismic observations and (bottom) tsunami importance  $I_t$ . Event type labels as in Figure 1.

does moment magnitude and that earthquakes with a high tsunamigenic potential (e.g., possible tsunami importance  $I_t \geq 2$  or  $I_t \geq 10$ ) can be rapidly and reliably identified through a procedure that determines if  $T_0$  is likely to exceed 50 or 100 s. This identification can be performed within 5–10 min after OT for most regions using currently available seismographic stations and probably in less than 3–5 min for regions with higher station density, such as Japan, Taiwan, Indonesia, the Mediterranean and Western North America. This identification forms a complement to initial estimates of the location, depth and magnitude of an earthquake to improve the reliability of tsunami early

warning, and, in some cases, may make possible such warning.

[16] **Acknowledgments.** We thank Alessio Piatanesi and two reviewers for helpful comments. This work is supported by the 2008–2010 Dipartimento della Protezione Civile S3 project. We use SeisGram2K (<http://www.alomax.net/software>) for seismogram analysis and figures and GMT (<http://gmt.soest.hawaii.edu/>) and OpenOffice.org Calc for graphs. The IRIS DMC (<http://www.iris.edu>) provided access to waveforms used in this study.

## References

- Bormann, P., and J. Saul (2009), Earthquake magnitude, in *Encyclopedia of Complexity and Systems Science*, edited by A. Meyers, Springer, Berlin, in press.
- Dziewonski, A., T. A. Chou, and J. H. Woodhouse (1981), Determination of earthquake source parameters from waveform data for studies of global and regional seismicity, *J. Geophys. Res.*, *86*, 2825–2852.
- Ekström, G., A. M. Dziewonski, N. N. Maternovskaya, and M. Nettles (2005), Global seismicity of 2003: Centroid-moment-tensor solutions for 1087 earthquakes, *Phys. Earth Planet. Inter.*, *148*, 327–351.
- Hara, T. (2007), Measurement of the duration of high-frequency energy radiation and its application to determination of the magnitudes of large shallow earthquakes, *Earth Planets Space*, *59*, 227–231.
- Hirshorn, B., and S. Weinstein (2009), Rapid estimates of earthquake source parameters for tsunami warning, in *Encyclopedia of Complexity and Systems Science*, edited by A. Meyers, Springer, Berlin, in press.
- Lomax, A. (2005), Rapid estimation of rupture extent for large earthquakes: Application to the 2004, M9 Sumatra-Andaman mega-thrust, *Geophys. Res. Lett.*, *32*, L10314, doi:10.1029/2005GL022437.
- Lomax, A., and A. Michelini (2005), Rapid determination of earthquake size for hazard warning, *Eos Trans. AGU*, *86*(21), 202.
- Lomax, A., and A. Michelini (2009),  $M_{wpd}$ : A duration-amplitude procedure for rapid determination of earthquake magnitude and tsunamigenic potential from P waveforms, *Geophys. J. Int.*, *176*, 200–214, doi:10.1111/j.1365-246X.2008.03974.x.
- Lomax, A., A. Michelini, and A. Piatanesi (2007), An energy-duration procedure for rapid determination of earthquake magnitude and tsunamigenic potential, *Geophys. J. Int.*, *170*, 1195–1209, doi:10.1111/j.1365-246X.2007.03469.x.
- Newman, A. V., and E. A. Okal (1998), Teleseismic estimates of radiated seismic energy: The  $E/M_0$  discriminant for tsunami earthquakes, *J. Geophys. Res.*, *103*, 26,885–26,898.
- Satake, K. (2002), Tsunamis, in *International Handbook of Earthquake and Engineering Seismology*, pp. 437–451, edited by W. H. K. Lee et al., Academic, Amsterdam.
- Tsuboi, S., K. Abe, K. Takano, and Y. Yamanaka (1995), Rapid determination of  $M_w$  from broadband P waveforms, *Bull. Seismol. Soc. Am.*, *85*, 606–613.
- Tsuboi, S., P. M. Whitmore, and T. J. Sokolowski (1999), Application of  $M_{wp}$  to deep and teleseismic earthquakes, *Bull. Seismol. Soc. Am.*, *89*, 1345–1351.

A. Lomax, ALomax Scientific, 161 Allee du Micocoulier, F-06370 Mouans-Sartoux, France. (anthony@alomax.net)

A. Michelini, Istituto Nazionale di Geofisica e Vulcanologia, Via di Vigna Murata 605, I-00143 Roma, Italy.